

# Statistical Analyses of Surface Temperatures in the IPCC Fifth Assessment Report

[Douglas J. Keenan](#)

17 December 2013    First version

28 December 2016    This version

Acknowledgements: for comments on drafts, I am grateful to [Sir David Cox](#), [David Henderson](#), [Demetris Koutsoyiannis](#), Janice Moore, and [William Nordhaus](#).

## 1. Introduction

In September 2013, the IPCC issued the first volume of its Fifth Assessment Report (AR5). The volume includes extensive discussions of observations of the average temperature on Earth's surface (i.e. where people live). The discussions include some statistical analyses of those observations. This critique considers the merits of such statistical analyses. No background in statistics is required.

## 2. How to do a statistical analysis

To understand statistical analysis, consider an example. Suppose that we toss two coins, and we ask what the probability is that both coins come up heads. To determine the probability, we will make two reasonable assumptions: (i) the probability that a coin comes up heads is  $\frac{1}{2}$  and (ii) one toss has no effect on the other toss. Then, the probability of two heads is calculated to be  $\frac{1}{2} \times \frac{1}{2} = \frac{1}{4}$ .

In general, any statistical analysis will consist of two phases. The first phase is to make some assumptions about the process that generates the data (in our example, we made two such assumptions). The second phase is to do some mathematical calculations (in our example, we did a simple multiplication).

The assumptions are obviously vital for the analysis. If, for instance, we had assumed that the probability a coin comes up heads is  $\frac{1}{3}$ , then our final answer would have been different:  $\frac{1}{9}$ . The assumptions, collectively, are called the "statistical model". Every statistical analysis thus vitally depends on what statistical model—i.e. set of assumptions—is chosen.

Although the mathematical calculation in our example was simple, it can sometimes be very complicated. Such complication used to be a major difficulty for statistical analyses (and if you studied statistics in the 20<sup>th</sup> century, you would have spent much effort considering how to do such complicated calculations). Nowadays, though, there is usually an easy way.

To illustrate the easy way, consider our example of tossing two coins. Instead of calculating the probability of getting two heads, we could estimate the probability as follows: take two coins, toss them a million times, and count the number of times that both coins come up heads. I tried doing that, and counted 249943 times that both coins came up heads; thus the estimated probability is  $249943/1000000$ , which is not exactly  $\frac{1}{4}$ , but is extremely close. The lack of exactness will almost always be negligible, in practice.

I did not actually toss coins a million times, of course. Rather, I used a computer program to roughly simulate doing that. The program was based on the two assumptions that we made. In other words, what I actually did was use a program that simulated the statistical model.


This method of running a program that simulates the model a large number of times can often be used, instead of doing the mathematical calculation. The method is called the “[Monte Carlo method](#)”. The method has been known for decades, but techniques for applying it quickly and accurately were first developed only around the year 2000. (The algorithm that led to this development is the [Mersenne Twister](#).)

The development of the Monte Carlo method implies that the second phase of a statistical analysis can often be done almost mechanically. Nowadays, then, it is often the first phase—selecting an appropriate model—that is the sole difficult task in statistical analyses.

### 3. Significant trends

Imagine tossing a coin ten times. If the coin came up Heads each time, we would have very good evidence that the coin was not a fair coin. Suppose instead that the coin was tossed only three times. If the coin came up Heads each time, we would not have good evidence that the coin was unfair: getting Heads three times can reasonably occur just by chance.

In Figure 1, each graph has three segments, one segment for each toss of a coin. If the coin came up Heads, then the segment slopes upward; if it came up Tails, then the segment slopes downward. In Figure 1, the graph on the left illustrates tossing Heads, Tails, Heads; the middle graph illustrates Tails, Heads, Tails; and the last graph illustrates Heads, Tails, Tails.



*Figure 1. Coin tosses: H, T, H (left); T, H, T (middle); H, T, T (right).*

Now consider Figure 2. At first, it might seem obvious that the graph shows an increase. This graph, however, illustrates Heads, Heads, Heads. Three Heads is not good evidence for anything other than random chance occurring. A statistician would say that although Figure 2 shows an increase, the increase is “not significant”.




Figure 2. Coin tosses: H, H, H.

Suppose that instead of tossing coins, we roll ordinary six-sided dice. If a die comes up 1, a line segment is drawn sloping downward; if it comes up 6, a segment is drawn sloping upward; and if it comes up 2, 3, 4, or 5, a segment is drawn straight across. We will roll each die three times. Some examples are given in Figure 3.




Figure 3. Dice rolls: 3, 6, 3 (left); 1, 5, 2 (middle); 4, 6, 1 (right).

Next consider Figure 4, which corresponds to rolling 6 three times. This outcome will occur by chance just once out of 216 times, and so it gives significant evidence that the die is not rolling randomly. That is, the increase shown in Figure 4 is significant.




Figure 4. Dice rolls: 6, 6, 6.

Note that Figure 2 and Figure 4 look identical. In Figure 2, the increase is not significant; yet in Figure 4, the increase is significant. These examples illustrate that we cannot determine whether a line shows a significant increase just by looking at it. Rather, we must know something about the process that generated the line.

In practice, we will not perfectly understand the process. What statisticians do instead is choose a model to represent the process as well as feasible.

An increase that is significant under one model might well be insignificant under another model, as illustrated. Put plainly, we can reach almost any conclusion, by (mis)choosing a suitable model. Again, the choice of model is vital.

#### 4. Time series

A concept from statistics that we need is that of a *time series*. A time series is any series of measurements taken at regular time intervals. Examples include the following: prices on the New York Stock Exchange at the close of each business day; the maximum temperature in London each day; the total wheat harvest in Canada each year. Another example is the average global temperature each year.

Suppose that today is extremely warm, at some location. Then there is a tendency for tomorrow to be warmer than average. More generally, what happens at some time in the future can be dependent upon what is happening now. Note that this is different from what happened when we tossed two coins: with the two coins, the outcome of the second toss was not dependent upon the outcome of the first toss.

Having future events be dependent upon the present complicates the statistical analysis of time series. Some elaboration on that is given in Excursus 1.

**Excursus 1.** As a first example, consider how the temperature today is correlated with the temperature tomorrow. The statistical analysis of daily temperatures should consider that correlation. Next, consider how the temperature during the past week will have some correlation with the temperature tomorrow. Again, the statistical analysis should consider that. Then consider how the temperature during the present season will tend to have some correlation with the temperature tomorrow. More generally, many different time scales are potentially relevant, and those time scales should be considered. Furthermore, the various correlations need not be linear. Properly accounting for all these issues is problematic.

The complications arising in the analysis of time series tend to make the selection of a statistical model for a time series difficult. Indeed, one of the world's leading specialists in time series, [Howell Tong](#), stated the following, in his book *Non-linear Time Series* (§5.4).

A fundamental difficulty in statistical analysis is the choice of an appropriate model. This is particularly pronounced in time series analysis.

The complications of time series also make developing intuition about time series difficult. For instance, it might be tempting to believe that we can determine whether an increase/decrease is likely to be due to chance (i.e. is not significant) just by looking at a plot of the data. Human intuition, however, can be misleading. For examples, some plots in Figure 5 might appear to display significant increases/decreases; yet all the plots were generated by running a purely-random statistical model of time series.





Figure 5. Some plots of random time series. (Plots were generated via a driftless ARIMA(3,1,0) model, with parameters set to the maximum likelihood estimates for HadCRUT 4.2.0.0 spanning 1880–2012.)

The plots in Figure 5 can be compared with the plot of global surface temperatures in Figure 6 (temperatures are shown relative to the 1961–1990 average).



*Figure 6. Global surface temperatures, 1880–2012. Temperatures are in °C, offset from the 1961–1990 mean. (Data from [HadCRUT 4.2.0.0.](#))*

In Figure 5, each plot has the same number of years, and the same starting temperature, as Figure 6. The plots in Figure 5 were generated using a model that has been suggested as a plausible candidate model for the series of global temperatures; the model is purely random. As shown, it is not unusual for the model to generate a time series that gives the illusion of having a trend that is at least as strong as the apparent trend in global temperatures.

The above again illustrates that, when determining whether an increase/decrease in a time series is significant, we cannot just look at a plot of the data. We must use statistical analysis.

## **5. Trend analysis in Chapter 2—Box 2.2**

In AR5, Volume I, Chapter 2 is devoted to observations of the atmosphere: observations of temperature, humidity, wind, etc. The statistical method used to evaluate trends in those observations is described in Box 2.2, which is subtitled “Trend models and estimation”. The first part of Box 2.2 is extracted below.

Many statistical methods exist for estimating trends in environmental time series (see Chandler and Scott, 2011 for a review). The assessment of long-term changes in historical climate data requires trend models that are transparent and reproducible, and that can provide credible uncertainty estimates.

Historical climate trends are frequently described and quantified by estimating the linear component of the change over time (e.g., AR4 [i.e. the IPCC Fourth Assessment

Report]). Such linear trend modelling has broad acceptance and understanding based on its frequent and widespread use in the published research assessed in this report, and its strengths and weaknesses are well known (von Storch and Zwiers, 1999; Wilks, 2006). Challenges exist in assessing the uncertainty in the trend and its dependence on the assumptions about the sampling distribution (Gaussian or otherwise), uncertainty in the data used, dependency models for the residuals about the trend line, and treating their serial correlation (Von Storch, 1999; Santer et al., 2008).

The quantification and visualisation of temporal changes are assessed in this chapter using a linear trend model that allows for first order autocorrelation in the residuals (Santer et al., 2008; Supplementary Material 2.SM.3). Trend slopes in such a model are the same as ordinary least squares trends; uncertainties are computed using an approximate method. The 90% confidence interval quoted is solely that arising from sampling uncertainty in estimating the trend.

There is no *a priori* physical reason why the long-term trend in climate should be linear in time. Climatic time series often have trends for which a straight line is not a good approximation (e.g., Seidel and Lanzante, 2004). The residuals from a linear fit in time often do not follow a simple autoregressive or moving average process, and linear trend estimates can easily change when estimates are recalculated using data covering shorter or longer time periods or when new data are added. When linear trends for two parts of a longer time series are calculated separately, the trends calculated for two shorter periods may be very different (even in sign) from the trend in the full period, if the time series exhibits significant nonlinear behavior in time (Box 2.2, Table 1).

The first paragraph is reasonable. The reference by Chandler and Scott is a book, which contains a section entitled “The linear trend” (§3.1); this section correctly states that “it is necessary to check the assumptions of any model before interpreting the results”. In other words, if the model’s assumptions have not been argued to be valid, then we cannot use the model to draw any inferences about the data.

The second paragraph correctly states that “Challenges exist in assessing the uncertainty in the trend”—and thus, in particular, assessing whether the trend is significant.

The third paragraph states that the IPCC has chosen a statistical model that comprises a straight line with first-order autocorrelated noise. If you are unfamiliar with such noise, that does not matter here. What is important here is that a model has been chosen, yet there is no scientific justification given for the choice. The failure to present any evidence or logic to support the assumptions of the model is a serious violation of basic scientific principles.



The fourth paragraph acknowledges that “residuals from a linear fit in time often do not follow a simple autoregressive ... process” (indeed, that is well known). This means that the chosen model does not fit the data; i.e. the model is acknowledged to be statistically inappropriate.

Box 2.2 concludes with this statement: “The linear trend fit is used in this chapter because it can be applied consistently to all the datasets, is relatively simple, transparent and easily comprehended, and is frequently used in the published research assessed here.”

Box 2.2 can be summarized as follows. A statistical model was chosen, without any statistical justification. Moreover, the chosen model is believed to be statistically inappropriate for climatic data. The model was chosen anyway for two reasons: first, choosing a more appropriate model would require more effort; second, almost everyone else has been using the same model—though also without statistical justification.

The first reason is plainly not a valid reason for choosing an inappropriate model. The Box is correct in stating that the chosen statistical model is “frequently used in the published research”; repeating an error many times, however, does not correct the error.

## **6. Trend analysis in Chapter 2—additional issues**

Chapter 2 also discusses statistics generally, in its Introduction (§2.1). Following is a relevant extract.

It is important to note that the question of whether the observed changes are outside the possible range of natural internal climate variability and consistent with the climate effects from changes in atmospheric composition is not addressed in this chapter, but rather in Chapter 10. No attempt has been undertaken to further describe and interpret the observed changes in terms of multidecadal oscillatory (or low frequency) variations, (long-term) persistence and/or secular trends (e.g., as in Cohn and Lins, 2005; Koutsoyiannis and Montanari, 2007; Zorita et al., 2008; Lennartz and Bunde, 2009; Mills, 2010; Mann, 2011; Wu et al., 2011; Zhou and Tung, 2012; Tung and Zhou, 2013). In this chapter, the robustness of the observed changes is assessed in relation to various sources of observational uncertainty (Box 2.1). In addition, the reported trend significance and statistical confidence intervals provide an indication of how large the observed trend is compared to the range of observed variability in a given aspect of the climate system (see Box 2.2 for a description of the statistical trend model applied).

This claim is key: “the question of whether the observed changes are outside the possible range of natural ... climate variability ... is not addressed in this chapter”. This chapter, though, does contain statistical analyses. The above text additionally claims that the purpose of those analyses is to “provide an indication of how large the observed trend is compared to the range of observed variability”.

In other words, the statistical analyses do not indicate whether the observed increases are outside the range of natural variability, but they do indicate if the observed increases are large compared to the range of variability. Obviously, the two claims conflict with each other.

Additionally, the term “observed trend” is misleading, because trends are generally meaningful only with confidence intervals (or similar). Confidence intervals, though, are not observed; rather, they are derived via the statistical model (which is chosen via human judgement).

Furthermore, by acknowledging that “No attempt has been undertaken to further describe and interpret the observed changes in terms of multidecadal oscillatory (or low frequency) variations, (long-term) persistence and/or secular trends ...”, Chapter 2 is (again) effectively acknowledging that there are statistical models that might well be more appropriate than the model that was chosen. And the chapter is deliberately avoiding considering those models.

The stated reason for not considering those models is that the purpose of the chapter is to consider “observational uncertainty”. For an example of observational uncertainty, we can examine the global surface temperature in 2012. The best estimate of that temperature is 0.45 °C (relative to the 1961–1990 average temperature). The value 0.45 °C, though, is not exact, because we do not have exact temperature measurements for every place in the world. Researchers, however, have stated that they are 90% confident that the exact temperature was in the interval 0.37–0.53 °C. That interval, known as a “90%-confidence interval”, is a way of indicating what the uncertainty is in the estimate of the 2012 temperature.

Chapter 2 is supposed to present observations of surface temperatures, and other aspects of the atmosphere, and to indicate how much uncertainty there is in each of those observations. Chapter 2 actually discusses uncertainty only a little.

Some illustration of the uncertainty for global temperatures is given in Figure 7. For the last year shown in the figure, 2012, there are two dots: one blue, at 0.37 °C; one red, at 0.53 °C. Thus, the dots indicate the 90%-confidence interval for the temperature in 2012. The other dots indicate similarly for the other years.




Figure 7. The 90%-confidence intervals for annual global surface temperatures, 1880–2012. For each year, the corresponding blue-red pair of dots denotes the end points of the confidence interval. Temperatures are in °C, offset from the 1961–1990 mean. (Data calculated from [HadCRUT 4.2.0.0.](#))

A straight line that was fit only via observational uncertainty would have to lie below almost all the red dots and above almost all the blue dots. Such a line obviously cannot exist. Hence, it is not possible to fit a straight line based only on observational uncertainty. Thus, the claim (in the chapter’s Introduction) that some trend line indicates observational uncertainty is false.

The main conclusion regarding Chapter 2 is as follows. Chapter 2 is not supposed to present statistical analyses, except to consider observational uncertainty; other statistical analyses are reserved for Chapter 10. Chapter 2 states that, but it presents trend analysis of the data anyway. The trend analysis is based on a model that is unjustified and that the chapter acknowledges is inappropriate.

☛ With AR5, overall responsibility for each chapter was assigned to “Coordinating Lead Authors”. In Volume I, Chapter 2 had three Coordinating Lead Authors ([Dennis L. Hartmann](#), [Albert M.G. Klein Tank](#), [Matilde Rusticucci](#)). The three were sent a draft of this critique, on 10 December 2013. I have not received a reply.

## 7. Trend analysis in Chapter 10

As noted above, statistical analysis of the series of global temperatures is officially within the remit of Chapter 10, not Chapter 2. The key issue for the statistical analysis is whether the increase in global temperatures is statistically significant. The IPCC refers to this issue as *detection*. Chapter 10 states the following: “An identified change is *detected* in observations if its likelihood of occurrence by chance due to internal variability alone is determined to be small” (§10.2.1, emphasis added). That is essentially the definition of *significant*.

The relevant section of Chapter 10 is §10.2.2, entitled “Time-series methods, causality and separating signal from noise”. The section begins by comparing the statistical analysis of data to the analysis that is done via supercomputer simulations of the global climate system. It states this: “The advantage of [time-series] approaches is that they do not depend on the accuracy of any complex global climate model, but they nevertheless have to assume some kind of model, or restricted class of models ...”. This statement is correct—and crucial.

The section also includes the following statements.

Time-series methods ultimately depend on the ... adequacy of the statistical model employed.

The assumptions of the statistical model employed can also influence results.

All these [time-series] approaches are subject to the issue of confounding factors....

Again, all of this is correct. In stating these things, the section is presenting the basics of the statistical situation reasonably fairly.

Additionally, §10.2.2 states this: “Trends that appear significant when tested against [the statistical model used in Chapter 2] may not be significant when tested against [some other statistical models]”. Thus, §10.2.2 effectively acknowledges that the statistical model used in Chapter 2 should not have been relied on.

So, what statistical model does §10.2.2 choose? None. That is, §10.2.2 effectively acknowledges that we do not understand the data well enough to choose a statistical model. It does that even though it also acknowledges that choosing such a model is required for drawing inferences. The conclusion is thus clear: it is currently not possible to draw inferences from the series of global temperatures. This conclusion is extremely important. It should have been stated explicitly, and it should have been noted in the Executive Summary of Chapter 10.

Although this critique is focused on surface temperature observations, the same statistical criticism applies to other claims of observational evidence for significant global warming. Simply put, no one has yet presented valid statistical analysis of any observational data to show global warming is real. Moreover, that applies to any warming—whether attributable to humans or to external natural factors, such as the sun. This is implied by §10.2.2, and indeed it is clear from the statistics.

## 8. Summary for Policymakers

The most important part of AR5 is arguably the [Summary for Policymakers](#) (SPM) in Volume I. The SPM synthesizes all the chapters in the volume, and it is intended to directly influence national governments.

The first section of the SPM, after the introduction, lists bullet points of evidence for global warming. The first bullet point thus appears as the single most important piece of evidence for global warming, from the perspective of policymakers. The first bullet point begins as follows.

The globally averaged combined land and ocean surface temperature data as calculated by a linear trend, show a warming of 0.85 [0.65 to 1.06] °C, over the period 1880 to 2012....

(The numbers in square brackets indicate the 90%-confidence interval. Having the confidence interval so far away from including 0 implies that the trend is extremely significant.)

The claim in the bullet point is essentially copied from the Executive Summary of Chapter 2, which claims that “global combined land and ocean surface temperature data show an increase of about 0.85 [0.65 to 1.06] over 1880–2012 ... when described by a linear trend”. The claim, however, is untenable, as discussed above and as acknowledged by §10.2.2.

Simply put, the SPM ignores what is said in Chapter 10. It does that even though responsibility for the statistical analysis lies with Chapter 10.

## 9. Trend analysis by the Met Office

The statistical model used in Chapter 2 of AR5, Volume I, was also used in Fourth Assessment Report (AR4), published in 2007. As a result of being used in AR4, the model has been studied by, amongst others, the Met Office, which is the primary institute for global-warming research in the UK. (The Met Office was originally known as the “Meteorological Office”.)

A research paper studying the statistical model used in Chapter 2 was published by the Met Office Chief Scientist, [Julia Slingo](#), in May 2013. The paper is entitled “[Statistical models and the global temperature record](#)”. It effectively acknowledges that the statistical model of Chapter 2 is untenable for the series of global temperatures. Moreover, although it considers other models, it does not attempt to select a model for the series. In short, the paper comes to essentially the same conclusion as §10.2.2.

The events that led to the writing of the paper are noteworthy. Briefly, [Lord Donoughue](#) submitted a Parliamentary Question that asked if the rise in global temperatures since 1880 was statistically significant. The Answer, which was sourced from the Met Office, was *yes*. The statistical model upon which the answer was based, however, was the statistical model used in AR4 (and Chapter 2 of AR5). Hence, I contacted Lord Donoughue, explained that the Answer was untenable, and offered my services as a statistical adviser.

Lord Donoughue then submitted another Parliamentary Question, which essentially asked if the statistical model used for the prior Question was justified. The Met Office refused to answer the Question. Lord Donoughue asked again, and again—five times in total—and he was refused each time.

The rules of Parliament require Parliamentary Questions to be answered. Lord Donoughue, though, was unsure about how to enforce that; indeed, he would later [remark](#), “In 28 years in Parliament I do not recall such obfuscation”. In consequence, Lord Donoughue consulted with the Leader of the House of Lords and the Deputy Clerk of the Parliaments. He then sent a letter to the responsible minister, the Parliamentary Under Secretary of State for Energy and Climate Change.

The minister, [Baroness Verma](#), had previously been signing off on answers obtained from the Met Office, on the assumption that the Met Office was acting in good faith. Upon receipt of the letter, the minister required the Met Office to answer the question properly. The proper answer was then given in Parliament: the statistical model is untenable. Moreover, the Met Office [told](#) Parliament that it did “not use a linear trend model to detect changes in global mean temperature”. The paper written by the Chief Scientist, cited above, is essentially an elaboration on that.

For more details on the foregoing events, see the Bishop Hill blog post entitled “[Met Office admits claims of significant temperature rise untenable](#)”. Simply put, the Met Office, in particular its Chief Scientist, was attempting to mislead Parliament.

There have been other situations where Chief Scientist Slingo made statistical claims about the climate that she knew were highly misleading. An example is described in the Bishop Hill blog post “[Climate correspondents](#)”.

In AR5, Volume I, the SPM relies upon the same statistical model as the IPCC had used in AR4—as discussed above. Hence, I sent the following message to Chief Scientist Slingo.

The IPCC’s AR5 WGI Summary for Policymakers includes the following statement.

The globally averaged combined land and ocean surface temperature data as calculated by a linear trend, show a warming of 0.85 [0.65 to 1.06] °C, over the period 1880–2012....

(The numbers in brackets indicate 90%-confidence intervals.) The statement is near the beginning of the first section after the Introduction; as such, it is especially prominent.

The confidence intervals are derived from a statistical model that comprises a straight line with AR(1) noise. As per your paper "[Statistical models and the global temperature record](#)" (May 2013), that statistical model is insupportable, and the confidence intervals should be much wider—perhaps even wide enough to include 0 °C.

It would seem to be an important part of the duty of the Chief Scientist of the Met Office to publicly inform UK policymakers that the statement is untenable and the truth is less alarming. I ask if you will be fulfilling that duty, and if not, why not.

I did not receive a reply. In an attempt to get an answer, Lord Donoughue has now submitted some further Parliamentary Questions.

## 10. The crucial question

The crucial question is this: *what statistical model should be chosen?* Both the IPCC (§10.2.2) and the Met Office have considered that question; neither found an answer. Indeed, finding an answer would require some original research.

The central issue here is simple, and does not require training in statistics to understand. The central issue is this: if assumptions are made in a scientific analysis, then those assumptions should not be merely proclaimed, but rather given some scientific justification. Yet, virtually all statistical analyses of climatic data proceed by merely proclaiming some assumptions.

The full situation is even worse, because the assumptions that are commonly used in statistical analyses of climatic data are not only unjustified, but also unjustifiable; i.e. it is known that the assumptions are inappropriate for the data. An example of that is in AR5, Volume I: the statistical analysis in the chapter on atmospheric observations (Chapter 2) relies on an assumption that is known to be inappropriate. Astoundingly, the inappropriateness is acknowledged in the chapter, as well as in another chapter which has the responsibility for the statistical analysis (Chapter 10).

There seems to be only one scientist who has seriously attempted to answer the crucial question, i.e. to choose a statistical model. That scientist is Demetris Koutsoyiannis, at the

National Technical University of Athens. Koutsoyiannis has not (yet) found a viable answer to the question; at least, though, he has tried to. No other researcher has tried, to my knowledge. For an introduction to some of the work of Koutsoyiannis, see Excursus 2.

**Excursus 2.** The number of statistical models that could potentially be considered for the global temperature series is infinite. How can we choose among those models? A leading statistician in the U.S. said the following, in an e-mail to me.

My sense is that the observed time series is not sufficiently long to cleanly distinguish among various time series models, nor to definitively demonstrate man-made warming versus natural cycles versus (for some models) a mostly flat trend.

Indeed, that should be obvious to anyone who has reasonable skill at the analysis of time series. It is only true, however, if we are considering purely-statistical analyses. Generally, though, analyses of data should incorporate some knowledge of the application area: in this case, the physics of the climate system. That is, we should try to use physics to constrain the set of candidate models. That strategy has also been [suggested](#) by a statistician at the Met Office, Doug McNeall.

Although that strategy is clear and arguably necessary, implementing it seems to be extremely difficult. The only researcher who has attempted implementing it, as far as I know, is Koutsoyiannis. Koutsoyiannis invokes thermodynamic constraints, in particular. For a non-technical overview of his approach, see the Bishop Hill blog post "[Koutsoyiannis 2011](#)".

The reliance on merely proclaimed assumptions, in statistical analyses of climatic data, implies that virtually all claims to have drawn statistical inferences from climatic data are untenable. In particular, there is no demonstrated observational evidence for significant global warming.

## 11. Conversations with some British climate scientists

I had an [e-mail discussion](#) with a Senior Scientist at the Met Office about the existence of evidence for global warming. The scientist, [Vicky Pope](#), had published an [article](#) about global warming in *The Guardian* (on 23 March 2012). The article claimed that “[global warming] is a matter of science and therefore of evidence – and there’s lots of it”, that a “whole range of different datasets and independent analyses show the world is warming”, and that there is



“overwhelming evidence for man-made climate change”. The article, however, did not substantiate those claims.

Hence, I e-mailed Pope, saying “I ask you to detail a single piece of observational evidence, and supporting analysis, for global warming”. Pope replied politely, but her reply did not specify any evidence. I answered, *inter alia*, “I note that your message does not present any piece of observational evidence, despite my asking for one piece”. Pope again replied politely, but again her reply did not specify any evidence; rather, her reply said “I will think about how and where to respond to the particular points that you raise”.

Thus, it seems that a Senior Scientist at the Met Office is aware that there is no demonstrated observational evidence for (significant) global warming. As noted earlier, it also seems that the Chief Scientist at the Met Office is aware. And now, with AR5, there seems to be awareness in Volume I, §10.2.2.

You might well ask how the misperception that observational evidence exists could have become widespread. Part of the problem is that climatologists generally have no training in the statistical analysis of time series—even though almost all climatic data sets are time series. In other words, climatologists do not know how to statistically analyse climatic data.

An exemplification of this occurred at a [debate](#) related to global warming that was held in July 2010. The debate was hosted by *The Guardian*. There were five panellists, including a former chair of the IPCC, [Bob Watson](#), and myself. Another one of the panellists was a Pro Vice Chancellor from the University of East Anglia, Trevor Davies; Davies oversees the work done at the university’s Climatic Research Unit (CRU), which is the most prominent institution for global-warming research in the UK after the Met Office; he was previously a researcher at CRU.

The current head of research at CRU is [Phil Jones](#), who is considered by many people to be the most eminent climatologist in the UK. Jones’ specialty is analysing climatic data. From my reading of some of Jones’ research publications, though, I concluded that Jones has negligible competence at data analysis. For that reason, I stated the following, during the debate.

I think people are really overestimating the competence and the skill of some of these scientists. Phil Jones, for example, could absolutely not pass an examination in an introductory undergraduate course in statistical time series.

After the debate, Davies told me that Jones could pass such an examination. I then offered to pay £500/minute to have Jones write the examination. My offer was not accepted. Thus, Davies seemed to implicitly admit that one of the world’s leading specialists in analysing climatic data did not have any competence at statistically analysing such data.

Jones, though, should not be singled out for criticism here. A lack of competence with time-series is exhibited by almost all climatologists (including those sceptical about global warming).

There have been some attempts to persuade climatologists to consider statistical analysis of time series more carefully, by both myself and others. Such attempts are almost always rebuffed. It is easy to criticize climatologists for that; before doing so, though, it is worth attempting to put yourself in their place.

Imagine that you had earned a Ph.D. in climatology: that takes about five years of hard work, on very tiny pay. Then you worked hard for decades more, earned respect from your peers, and essentially founded your professional identity on being an expert in the study of the climate system. And now, someone comes along and tells you that most of the work you and your colleagues have done during your careers is invalid, due to a statistical problem. How would you respond? Would you say, “Oh, that’s nice—thanks for letting me know”?

Scientists are human, and they respond in human ways. One key to understanding what has happened with climate science is to consider not just the science, but also the scientists.

## 12. The Australian experience

The Australian government commissioned a study on the impacts of global warming on the Australian economy. The study was led by a distinguished professor of economics in Australia, [Ross Garnaut](#), and it is known as the “[Garnaut Review](#)”. The Review began by assessing the scientific basis for global warming. That assessment concluded that global warming is a serious threat to the world.

Garnaut conducted the assessment of the scientific basis as follows. First, he recognized that almost all climatic data sets are time series, and thus analysing climatic data requires doing time-series analysis. Second, he commissioned two time-series specialists in Australia to analyse the series of global surface temperatures. Third, he founded the Review, in part, on the results of that analysis. In short, the scientific foundation of the Review was conducted in an exemplary way.

The statistical analysis that served as the foundation of the Garnaut Review is detailed in a paper, “[Global temperature trends](#)”, written by the two time-series specialists—[Trevor Breusch](#) and Farshid Vahid. I found invalidating errors in that analysis. I e-mailed Breusch and Vahid about the errors in June 2011. Breusch replied politely, but seemed to not understand the issues. I responded, elaborating and giving references. There were no further e-mails.

The statistical errors are actually clear and basic. A slightly oversimplified description of the errors is given in Excursus 3.

**Excursus 3.** When choosing a statistical model, one of the methods used by statisticians is to employ what is called “relative likelihood”. Using this method, we can tell that one model is, e.g., 1000 times more likely than another model to be the better model of the data (where “better” is defined in a certain technical way).

If one model is 1000 times more likely than another, second, model, then we would conclude that the second model should be rejected. If the first model is only a few times more likely than the second model, though, then we should not reject the second model. The situation here is analogous to gambling: if the odds are 1000 to 1 in our favour, then we are almost certain to win (in the analogy, be choosing the better model); if the odds are only 3 to 1 in our favour, then we might well lose (in the analogy, be choosing the worse model).

Breusch & Vahid considered some statistical models for the temperature series. The models that they considered, though, were all linear. Excluding nonlinear models is questionable. Indeed, the IPCC has previously noted that “we are dealing with a coupled non-linear chaotic system” (AR3, Volume I, [§14.2.2.2](#)).

Having restricted their consideration to only linear models, Breusch & Vahid then chose a model for which the increase in global temperatures is significant. Yet there were other models that were nearly as likely as the chosen model: thus, the choice of Breusch & Vahid was ill founded.

With some of the other models that were nearly as likely as the chosen model, the increase in temperatures is not significant. To summarize—some likely model shows the increase as significant and other likely models show the increase as not significant. Hence, we cannot determine whether the increase is significant. The main conclusion of Breusch & Vahid, however—based on their choice of model—is that the increase is significant. The main conclusion is thus actually baseless.

It is notable that the Met Office Chief Scientist, in her paper cited above, considered some linear models very similar to those considered by Breusch & Vahid. The Chief Scientist reasoned that the likelihood comparisons were “inconclusive”; so she did not choose among the models. She did find that the most likely model was one for which the increase in temperatures is significant. She noted, however, that there were other models that were nearly as likely and for which the increase is not significant. Hence, she drew no inferences regarding significance. In other words, when in essentially the same situation as Breusch &

Vahid, the Chief Scientist reached the correct conclusion. (This conclusion is also effectively implied by the more general considerations of AR5, Volume I, §10.2.2.)

☛ Breusch and Vahid were sent a draft of this critique, on 27 October 2013. I have not received a reply.

### **13. An example from outside climatology**

Vital problems with statistical analyses exist in fields of research other than global-warming science. An example from another field is described here. The field is [tephrochronology](#), which studies the chemistry of volcanic ash.

In statistics, there is a concept known as “standard error”; the concept is taught in all introductory statistics courses. A substantial portion of modern tephrochronology, though, is substantially based on misinterpreting the concept. The misinterpretation has sometimes led to conclusions that are the opposite of what would be concluded if the correct interpretation were used. I published a [paper](#) on this, in 2003. My paper was published in the leading journal for geochemistry, but it has since been largely ignored.

Not all fields of science do statistical analyses incompetently though. One branch of science where gross statistical incompetence occurs only rarely is medical science. Up until this century, however, statistical analyses in medical science were often appalling. A statistician at Oxford University, [Doug Altman](#), campaigned against incompetent statistics in medical research. His campaign included publishing works such as “[The scandal of poor medical research](#)” and “[Poor-quality medical research](#)”. After about a decade of this, the International Committee of Medical Journal Editors agreed to a major change in the way research manuscripts are peer reviewed: the Committee’s [Uniform Requirements for Manuscripts](#) was changed to include strong requirements for statistical methods. Moreover, if a manuscript relies on non-trivial statistics, medical journal editors now commonly call in a statistician as an extra reviewer. Since the change, statistical analyses in medical science have dramatically improved (though they are still far from perfect).

The foregoing illustrates that vital problems with statistical analyses exist, or have existed, in scientific fields other than climatology. Such problems can be at the level of an introductory statistics course, as in tephrochronology. Such problems can have enormous consequences, as in medicine. And rectification of such problems can be fought against by almost all scientists in the relevant field, as in both tephrochronology and medicine. Any attempt to understand the source of the problems in global-warming science should consider that.

## Appendix. Foundations of statistics

The main foundation of climatology is agreed upon by all climatologists: it comprises Newtonian mechanics, classical chemistry, etc. The main foundation of statistics, though, is not agreed upon by statisticians: rather, there are different contenders for the foundation. One of those contenders is called *frequentism*. The frequentist foundation inherently leads to using significance levels, confidence intervals, etc.; it is the best known foundation among climatologists, by far. Because frequentist statistics is so well known among climatologists, it has been generally adopted in this critique. The issues raised in this critique, though, are not dependent upon the foundation: all the issues can be recast for other foundations (in particular, for the Bayesian foundation and for the information-theoretic foundation).

---

---

The reason for so much bad science is not that talent is rare, not at all; what is rare is character. People are not honest, they don't admit their ignorance, and that is why they write such nonsense.

—*Sigmund Freud*

Half the harm that is done in this world is due to people who want to feel important. They don't mean to do harm—but the harm does not interest them. Or they do not see it, or they justify it because they are absorbed in the endless struggle to think well of themselves.

—*T.S. Eliot*

You should, in science, believe logic and arguments, carefully drawn, and not authorities.

—*Richard P. Feynman*